


This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Original) Process for the production of C₁-C₁₅-epothilone fragments of general formula I,



in which

R^{1a}, R^{1b} are the same or different and mean hydrogen, C₁-C₁₀-alkyl, aryl,

C₇-C₂₀-aralkyl, or together mean a -(CH₂)_m group with m = 2, 3, 4 or 5,

R^{2a}, R^{2b} are the same or different and mean hydrogen, C₁-C₁₀-alkyl,

C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, aryl, C₇-C₂₀-aralkyl or together mean a -(CH₂)_n group with n = 2, 3, 4 or 5,

R³ means hydrogen, C₁-C₁₀-alkyl, aryl, C₇-C₂₀-aralkyl,

R^{4a}, R^{4b} are the same or different and mean hydrogen, C₁-C₁₀-alkyl, aryl,

C₇-C₂₀-aralkyl or together mean a -(CH₂)_p group with p = 2, 3, 4 or 5,

R⁵ means hydrogen, C₁-C₁₀-alkyl, aryl, C₇-C₂₀-aralkyl,

R⁶, R⁷ each mean a hydrogen atom, together an additional bond or together an

oxygen atom,

G means a group $X=CR^8$ -, a bicyclic or tricyclic aryl radical,

R^8 means hydrogen, halogen, C_1 - C_{20} -alkyl, aryl, C_7 - C_{20} -aralkyl, which all can be substituted,

X means an oxygen atom, two alkoxy groups OR^{23} , a C_2 - C_{10} -alkylene- α,ω -dioxy group, which can be straight-chain or branched, H/OR^9 or a grouping $CR^{10}R^{11}$,

whereby

R^{23} stands for a C_1 - C_{20} -alkyl radical,

R^9 stands for hydrogen or a protective group PG^X ,

R^{10} , R^{11} are the same or different and stand for hydrogen, a C_1 - C_{20} -alkyl, aryl, or C_7 - C_{20} -aralkyl radical, or R^{10} and R^{11} together with the methylene carbon atom together stand for a 5- to 7-membered carbocyclic ring,

R^{13} means CH_2OR^{13a} , CH_2 -Hal, CHO, CO_2R^{13b} , or COHal,

R^{14} means hydrogen, OR^{14a} , Hal, or OSO_2R^{14b} ,

R^{13a} , R^{14a} mean hydrogen, SO_2 -alkyl, SO_2 -aryl, SO_2 -aralkyl or together a $-(CH_2)_o$ group or together a $CR^{15a}R^{15b}$ group,

R^{13b} , R^{14b} mean hydrogen, C_1 - C_{20} -alkyl, aryl, C_1 - C_{20} -aralkyl,

R^{15a} , R^{15b} are the same or different and mean hydrogen, C_1 - C_{10} -alkyl, aryl, C_7 - C_{20} -aralkyl, or together a $-(CH_2)_q$ group,

o means 2 to 4,

q means 3 to 6,

R²⁰ means OPG³, NHR²⁹, or N₃,

Z means an oxygen atom or H/OR¹²,


whereby

R¹² is hydrogen or a protective group PG^Z

including all stereoisomers as well as mixtures thereof, and

free hydroxyl groups in R¹³ and R¹⁴ can be etherified or esterified, free carbonyl groups in Z and R¹³ can be ketalized, converted into an enol ether or reduced, and free acid groups in R¹³ and R¹⁴ can be converted into their salts with bases, characterized in that

a C1-C6 fragment (epothilone numbering system) of general formula A




in which

R^{1a'}, R^{1b'}, R^{2a'}, R^{2b'}, R^{13'} and R^{14'} have the meanings already mentioned for R^{1a}, R^{1b}, R^{2a}, R^{2b}, R¹³ and R¹⁴, including all stereoisomers as well as mixtures thereof, and free hydroxyl groups in R¹³ and R¹⁴ can be etherified or esterified, free carbonyl groups in A and R¹³ can be ketalized, converted into an enol ether or reduced, and free acid groups in A

can be converted into their salts with bases,

is reacted with a C7-C12 fragment (epothilone numbering system) of general formula



in which

$R^{3a'}$, $R^{4a'}$, $R^{4b'}$ and $R^{5'}$ have the meanings already mentioned for R^{3a} , R^4 and R^5 , and


V means an oxygen atom, two alkoxy groups OR^{17} , a C_2 - C_{10} -alkylene- α,ω -dioxy group, which can be straight-chain or branched, or H/OR^{16} ,

W means an oxygen atom, two alkoxy groups OR^{19} , a C_2 - C_{10} -alkylene- α,ω -dioxy group, which can be straight-chain or branched, or H/OR^{18} ,

R^{16} , R^{18} , independently of one another, mean hydrogen or a protective group PG^1 ,

R^{17} , R^{19} , independently of one another, mean C_1 - C_{20} -alkyl,

to form a partial fragment of general formula AB




in which

$R^{1a'}$, $R^{1b'}$, $R^{2a'}$, $R^{2b'}$, $R^{3'}$, $R^{4a'}$, $R^{4b'}$, R^5 , $R^{13'}$, $R^{14'}$, V and Z have the already-mentioned meanings, and

PG^{14} represents a hydrogen atom or a protective group PG, and

this partial fragment of general formula AB is reacted with a C13-C15 fragment (epothilone numbering system) of general formula C



in which

G' has the meaning already mentioned in general formula I for G, and

$R^{7'}$ means a hydrogen atom,


$R^{20'}$ means halogen, N_3 , NHR^{29} , a hydroxy group, a protected hydroxy group O- PG^3 , a protected amino group $NR^{29}PG^3$, a C_1 - C_{10} -alkylsulfonyloxy group, which optionally can be perfluorinated, a benzoyloxy group that is optionally substituted by C_1 - C_4 -alkyl, nitro, chlorine or bromine, an

NR²⁹SO₂CH₃ group, an NR²⁹C(=O)CH₃ group, or a CH₂-C(=O)-CH₃ group,

R²¹ means a hydroxy group, halogen, a protected hydroxy group OPG³, a phosphonium halide radical PPh₃⁺Hal⁻ (Ph = Phenyl; Hal = F, Cl, Br, I), a phosphonate radical P(O)(OQ)₂ (Q = C₁-C₁₀-alkyl or phenyl) or a phosphine oxide radical P(O)Ph₂ (Ph = Phenyl),

R²⁹ means hydrogen or C₁-C₆-alkyl,

to form a compound of general formula ABC (= compound of general formula I)



in which

R^{1a'}, R^{1b'}, R^{2a'}, R^{2b'}, R^{3'}, R^{4a'}, R^{4b'}, R^{5'}, R⁶, R⁷, R¹³, R¹⁴, G and Z have the already mentioned meanings, and

PG¹⁴ represents a hydrogen atom or a protective group PG.

2. (Original) Process according to claim 1, wherein a compound of general formula

I,

in which

R^{1a}, R^{1b} are the same and mean C₁-C₆-alkyl, or together mean a -(CH₂)_m group

with m = 2, 3 or 4,

R^{2a}, R^{2b} are different and mean hydrogen, C₁-C₆-alkyl, C₂-C₁₀-alkenyl,

C₂-C₁₀-alkynyl or C₇-C₂₀-aralkyl,

R⁵ means hydrogen, C₁-C₆-alkyl,

R⁸ means hydrogen, halogen, C₁-C₆-alkyl,

R^{15a}, R^{15b} are the same or different and mean hydrogen, C₁-C₆-alkyl, aryl, C₇-

C₂₀-aralkyl, or together mean a -(CH₂)_q group,

q means 3 to 6,

is produced.

3. (Original) Process according to claim 1, wherein a compound of general formula

I,

in which

R^{1a}, R^{1b} are the same and mean C₁-C₃-alkyl, or together mean a -(CH₂)_m group

with m = 2, 3 or 4,

R^{2a} means hydrogen,

R^{2b} means C₁-C₅-alkyl, C₂-C₆-alkenyl, or C₂-C₆-alkynyl,

R⁵ means hydrogen, or C₁-C₃-alkyl,

R⁶, R⁷ together mean an additional bond,

G means a group X=CR⁸-, or a bicyclic aryl radical,

R⁸ means hydrogen, fluorine, chlorine, or C₁-C₃-alkyl,

X means oxygen or a group CR¹⁰R¹¹,

R¹⁰ means hydrogen,

R¹¹ means aryl,

R¹³ means CH₂OR^{13a} or CO₂R^{13b},

R¹⁴ means OR^{14a},

R^{13a}, R^{14a} together mean a CR^{15a}R^{15b} group,

R^{13b} means hydrogen or C₁-C₆-alkyl,

R^{15a}, R^{15b} are the same and mean C₁-C₃-alkyl, or together mean a -(CH₂)_q

group, or


R^{15a}, R^{15b} are different and mean hydrogen or aryl,

q means 4 or 5,

Z means oxygen,

is produced.


4. (Currently Amended) Process for the production of epothilone derivatives of general formula II



in which substituents R^{1a} , R^{1b} , R^{2a} , R^{2b} , R^3 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 , G , OPG^2 and Z have the meanings that are indicated in general formula I, and

A–K means a group $-O-C(=O)-$, $-OCH_2-$, $-CH_2C(=O)-$, $-NR^{29}-C(=O)-$, or $-NR^{29}-SO_2-$, wherein an initial epothilone product of general formula I that is obtained according to claim 1 ~~one of the preceding claims 1 to 3~~ is cyclized.

5. (Original) Compounds of general formula AB



in which $R^{1a'}$, $R^{1b'}$, $R^{2a'}$, $R^{2b'}$, $R^{3'}$, $R^{4a'}$, $R^{4b'}$, R^5 , $R^{13'}$, $R^{14'}$, V and Z have the already mentioned meanings, and PG^{14} represents a hydrogen atom or a protective group PG.